



SONOMA COUNTY VITICULTURE NEWSLETTER



December 2000

Rhonda J. Smith, Viticulture Farm Advisor

Save the date - Sonoma County Grape Day – Friday, February 16, 2001

The topics to be presented include results from a current Central Coast trellising trial in Chardonnay and Merlot, an update on latent grapevine viruses, and biological control of glassy-winged sharpshooters. Grape Day registration materials will be mailed in January. It will be held at the Luther Burbank Center in Santa Rosa.

New Publication Available:

“Riparian Vegetation Management for Pierce’s Disease in North Coast California Vineyards”

This manual describes a method to reduce the incidence of Pierce’s disease in vineyards adjacent to riparian areas. Managing riparian vegetation can reduce the numbers of blue-green sharpshooters – the principle insect vector of the bacterium that causes Pierce’s disease in the North Coast. It will give growers the information they need to effectively and legally modify and manage the riparian vegetation immediately adjacent to the edge of vineyard blocks that are prone to Pierce’s disease infections.

The manual describes an overview of selective host plant removal and revegetation methods that includes the key components of the information required in the Streambed Alteration Agreement Application that must be submitted to the California Department of Fish and Game. It provides information that can help the grower accomplish the goal of reducing the incidence of Pierce’s disease while simultaneously protecting fish and wildlife habitat.

This manual was written and reviewed by the Pierce’s disease/Riparian Habitat Workgroup, an informal

group of growers, resource and regulatory agencies, UC faculty and Cooperative Extension advisors and vegetation restoration specialists located in Sonoma and Napa Counties. It was coordinated by Ellie Insley. The Workgroup received funding from the North Coast Pierce’s Disease Task Force and the USDA, Natural Resource Conservation Service – Environmental Quality Incentives Program. In addition, Clos du Bois Winery, Robert Mondavi Winery, Sonoma County Grape Growers Association and the Sonoma County Viticulture Technical Group funded printing.

The manual is available on the web at the following address:

https://nature.berkeley.edu/xylella/pd_management/pd_north_coast/

A limited number of printed copies are available free of charge at UC Cooperative Extension offices in Sonoma, Napa and Mendocino Counties. Once my office is out of printed copies, photocopies will be provided and photocopying costs may apply.

UC Cooperative Extension offices
Sonoma County (707) 365-2621
Napa County (707) 253-4221
Mendocino County (707) 463-4495

What, No Surflan! What Do I Do?

Kurt Hembree
Fresno County Farm Advisor

If you are a grower of tree or vine crops in California, I'm sure you are aware by now that Surflan production for the 2001 season will be significantly impacted. According to Dow AgroSciences, a recent explosion near their southern manufacturing facility has halted production of one of the more popular tree and vine preemergence herbicides. As a result, amounts available to growers will be *extremely limited* this coming season. Therefore, your current stock of Surflan herbicide will most likely be all that is available. Surflan has been very effective at controlling a wide-array of annual grasses (like annual bluegrass, barnyardgrass, foxtails, etc.) and a select number of broadleaves (like chickweed, lambsquarters, purslane, and pigweeds). It is routinely tank-mixed with a broadleaf preemergence herbicide (usually Goal) to provide broad-spectrum weed control. Single or split applications are often made to obtain season-long control.

While it would be nice to say that there are plenty of alternatives and you have nothing to worry about, this is obviously not the case. Because growers are left with limited options, it will be very important that growers carefully select the most appropriate course of action to control their weeds. Certain considerations should be taken when deciding on which alternatives are the "best".

The best starting point is to have a full understanding of which weeds you expect to have in your field. If you have a whole host of weeds that require something other than Surflan for control, this is a good opportunity to look towards other options. If you are unsure of your weed spectrum, start looking for winter annuals as the temperatures begin to decline in the fall and the summer annuals in late winter or early spring.

While preemergence herbicides will not control the emerged weeds, it will give you a better understanding of what weeds you do have. You still have the option at that point of controlling them with a timely postemergence treatment (Roundup Ultra, Gramoxone, Touchdown, etc.).

Once you do know which weeds you are battling, you can select from the list of currently

registered pre- and/or postemergence herbicides that best fits your needs. The herbicide labels are pretty clear as to the specific weeds each will control. If unsure about the identity of certain weeds, contact your local farm advisor, PCA, or manufacturer representative for assistance. In many instances, you may have to use combinations of pre- and postemergence treatments to meet your goals. See table 1 for a list of herbicides that are registered in selected tree and vine crops in California.

Of the herbicides currently labeled, Devrinol or Solicam may offer the closest control when compared to Surflan and are registered in a majority of the tree and vine crops. However, it is important to understand that they will not work for all weed types or field conditions. For example, Devrinol must be incorporated within 4 to 7 days or the efficacy will decline. Time the treatment just ahead of a storm with a predicted rainfall of at least ½". Leasing additional spray equipment to get the job done on time or using a custom applicator may be needed where large acreage is a concern. Keep in mind, Devrinol tends to break down rapidly under wet conditions, especially in the spring and summer under frequent low-volume irrigation. Follow-up sprays with postemergence herbicides may also be required. In addition to the weeds Surflan controls, Devrinol will also control cudweed, filaree, prickly lettuce, pineappleweed, sowthistle, and others. Devrinol can be tank-mixed with other pre- and postemergence herbicides for broader weed control.

Solicam is perhaps a little weaker on the annual grasses, compared to Surflan, but is very effective at suppressing nutsedge. It is also effective on the mustard species, cudweed, and nightshade and will suppress filaree, groundsel, horseweed, and flaxleaved fleabane. Solicam can be tank-mixed with other pre- and postemergence herbicides for broader weed control. One of the major weaknesses of Solicam is its performance under light soil conditions and frequent low-volume irrigation. Solicam is fairly soluble in water and can be readily leached into the root zone, causing bleaching of the desirable crop foliage. Caution must be used to avoid possible phytotoxicity. It is important to adjust the rate based

on your particular soil texture. Please consult the appropriate label for recommended rates under varied soil types.

Finally, we do have a good selection of contact and systemic postemergence herbicides available. They are very effective at controlling most of the annual weeds you will encounter, but must be applied in a timely manner to keep them under control. If you choose to move towards a more postemergence approach to weed control, keep in mind that the weeds should be treated before they become hairy or mealy or prior to setting seed. For flaxleaved fleabane or horseweed control, it is best to treat when they are no more than 4” tall and before they begin to bolt. Some weeds like spurge can produce a large amount of seed rapidly (in less than 20 days after emergence), so timing of postemergence

sprays is critical to reduce the future population. Additional sprays or spot treatments will likely be needed near sprinkler emitters where weed flushes tend to be great. If entering wet fields is not an option for you in the winter, consider one of the preemergence herbicides, followed by timely postemergence treatments as access becomes available.

Several people are pursuing a Section 18 for the use of Visor in almonds and grapes. It is not known at this time if it will be approved, but if approved, will not likely be available until after the first of the year. So in the meantime, look at the other options as a possible solution to the Surflan shortage. If in doubt of your options, contact your local farm advisor, PCA or industry representative for assistance.

Table 1. Herbicide Registration Status in Selected Crops in California

Herbicides	Tree and Vine Crops											
	Almond	Apricot	Citrus	Fig	Grape	Kiwi	Nectarine	Olive	Peach	Pistachio	Plum	Walnut
<i>Preemerg.</i>												
Devrinol	R	R	R	R	R	R	R	R	R	R	R	R
Eptam	R	--	R	--	--	--	--	--	--	--	--	R
Gallery	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB
Goal	R	R	NB	R	R	R	R	R	R	R	R	R
Hyvar X	--	--	4+	--	--	--	--	--	--	--	--	--
Karmex	--	--	1+	--	3+	--	--	1+	3+	--	--	R
Kerb	--	--	--	--	1+	--	1+	--	1+	--	1+	--
Krovar	--	--	4+	--	--	--	--	--	--	--	--	--
Princep	1+**	--	1+	--	1+	--	1+	1+	1+	--	--	1+
Prowl	NB	NB	NB	--	NB	--	NB	--	NB	NB	NB	NB
Solicam	1.5+	1.5+	R	--	2+	--	1.5+	--	1.5+	--	1.5+	1.5+
Surflan	R	R	R	R	R	R	R	R	R	R	R	R
Treflan	R	--	R	--	R	--	R	--	R	--	R	R
Visor	NB	--	R	--	NB	--	--	--	--	NB	--	NB
<i>Postemerge.</i>												
Fusilade Dx	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB
Gramoxone Extra	R	R	R	R	R	R	R	R	R	R	R	R
MSMA	NB	NB	NB	--	NB	--	--	--	NB	--	NB	NB
Poast	R	NB	R	NB	R	NB	NB	NB	NB	NB	NB	R
Prism	NB	--	NB	--	NB	--	NB	NB	NB	NB	NB	NB
Roundup Ultra	R	R	R	R	R	R	R	R	R	R	R	R
Touchdown	R	R	R	NB	R	NB	NB	NB	R	NB	R	R
2,4-D	R	R	R	--	R	--	R	R	R	R	R	R

R = registered, -- = not registered, NB = non-bearing only, and + = number years must be established
 **Do not use on Mission variety

All herbicides mentioned in this article and chart are registered trade names and no endorsement of any particular herbicide is intended. This is not a recommendation for the use of herbicides. Please refer to the appropriate labels for application recommendations. 10/03/00

Sonoma County Viticulture Newsletter – December 2000

- Sonoma County Grape Day is February 16, 2001!
- New publication available.
- No Surflan?



Rhonda J. Smith
Viticulture Farm Advisor



The University of California prohibits discrimination against or harassment of any person employed by or seeking employment with the University on the basis of race, color, national origin, religion, sex, physical or mental disability, medical condition (cancer-related or genetic characteristics), ancestry, marital status, age, sexual orientation, citizenship, or status as a covered veteran (special disabled veteran, Vietnam-era veteran or any other veteran who served on active duty during a war or in a campaign or expedition for which a campaign badge has been authorized). University Policy is intended to be consistent with the provisions of applicable State and Federal laws. Inquiries regarding the University's nondiscrimination policies may be directed to the Affirmative Action/Staff Personnel Services Director, University of California, Agriculture and Natural Resources, 1111 Franklin, 6th Floor, Oakland, CA 94607-5200 (510) 987-0096. University of California, USDA, County of Sonoma Cooperating

ADDRESS SERVICE REQUESTED

49

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE

UC COOPERATIVE EXTENSION
2604 VENTURA AVE ROOM 100
SANTA ROSA CA 95403

Bulk Rate
US Postage Paid
USDA
Permit No. G-00268